

Study of the CN($^2\Sigma$) + N(4S) reaction at high temperatures

Otoniel Denis-Alpizar, and Rodrigo Urzúa-Leiva

Universidad Autónoma de Chile, Av. Pedro de Valdivia 425, Providencia, Santiago, Chile
otoniel.denis@uautonoma.cl

The CN molecule plays an important role in the chemistry around the surface of a spacecraft during the atmospheric re-entry. The collision of CN with others molecules and atoms has a particular interest in aerothermodynamic modeling.¹ This work focuses on studying the $CN+N \leftrightarrow N_2+C$ reaction within the triplet manifold $3A''$ of CN_2 . Recent theoretical studies have explored this reaction². However, the number of analytical potential energy surfaces (PES) is reduced, and computed at the CASPT3 level³ or limited to a two-dimensional surface⁴. Here, we developed a full dimensional high-level PES for this system from *ab initio* calculations at the MRCI-F12 level of theory. Then this new surface is used in quasi-classical trajectory calculations. Thermal rate coefficients are computed from 100 to 20000 K. The computed rates for the $CN+N \rightarrow N_2+C$ reaction are compared with available experimental data, and a good agreement is found. At low and intermediate temperatures, the N_2 formation is more efficient than the N-exchange, while at high temperatures, the rates for both channels are comparable. Finally, analytical modified Arrhenius expressions of the rates for the N_2 formation and N-exchange reactions are reported.

References

1. Martin, A., Cozmuta, I., Wright, M. J., & Boyd, I. D., *J. Thermophys. Heat. Trans.*, 2015, **29**, 222
2. Papakondylis, A., & Mavridis, A., *J. Phys. Chem. A*, 2019, **123**, 10290.
3. Frankcombe, T. J., McNeil, S. D., & Nyman, G., *Chem. Phys. Lett.*, 2011, **514**, 40
4. Ma, J., Guo, H., & Dawes, R., *Phys. Chem. Chem. Phys.*, 2012, **14**, 12090